

CERTIFICATION FORM: EpiPen® (epinephrine injection, USP) Auto-Injector for *EpiPen4Schools*® Replenishment Offer

Mylan will replenish your school's supply of EpiPen[®] (epinephrine injection, USP) or EpiPen Jr[®] (epinephrine injection, USP) Auto-Injectors prior to your annual eligibility date at no additional cost, provided that your school used the EpiPen[®] or EpiPen Jr[®] Auto-Injectors for authorized generic for EpiPen[®] or EpiPen Jr[®] Auto-Injectors for *EpiPen4Schools*[®] free product to treat a life-threatening allergic reaction (anaphylaxis) in your school.

First, please complete all of the fields below.

School Name/District Name:
School Address:
City/State/Zip:
School Phone:
School Contact Name:
School Contact Email:
What was the date of the anaphylactic event? MM/DD/YYYY
Where did the anaphylactic event occur? Class room Cafeteria Playground Gym Other
What was the suspected cause of the anaphylactic event? Food Insect Sting/Bite Latex Medication Other
 Did the person who experienced anaphylaxis have a known life-threatening allergy? Yes No Unknown
Was the person who experienced anaphylaxis a: Student Staff member Visitor Other
 Which epinephrine auto-injector was administered to treat the anaphylactic event? Authorized generic for EpiPen[®] Auto-Injector Authorized generic for EpiPen Jr[®] Auto-Injector EpiPen[®] Auto-Injector EpiPen Jr[®] Auto-Injector
Was more than one EpiPen [®] or EpiPen Jr [®] Auto-Injector or authorized generic for EpiPen [®] or EpiPen Jr [®] Auto-Injector administered to treat the anaphylactic event? • Yes • No
Was 911 or emergency medical services called? • Yes • No • Unknown
 Did the person who experienced anaphylaxis receive emergency medical care? Yes No Unknown

 Who administered the EpiPen[®] or EpiPen Jr[®] Auto-Injector or authorized generic for EpiPen[®] or EpiPen Jr[®] Auto-Injector to the person experiencing anaphylaxis (please do not name people directly)? School Nurse Student other than one experiencing the event Staff member Visitor
Self-Administered
Was the person who experienced anaphylaxis taken to the hospital?
• Yes
• No
Unknown
Is epinephrine your school's first-line treatment when an anaphylactic reaction occurs?
• Yes
• No
Unknown
If school policies permit, would you be willing to speak with us about the event?
• Yes
• No
I confirm that the EpiPen [®] or EpiPen Jr [®] Auto-Injector or authorized generic for EpiPen [®] or EpiPen Jr [®] Auto- Injector used in the anaphylactic event described above was received through the <i>EpiPen4Schools</i> [®] program.

Signature:

Date:

I certify that the information above is true and accurate.

Important Safety Information

Use EpiPen[®] (epinephrine injection, USP) 0.3 mg or EpiPen Jr[®] (epinephrine injection, USP) 0.15 mg Auto-Injectors right away when you have an allergic emergency (anaphylaxis). **Get emergency medical help right away**. You may need further medical attention. Only a healthcare professional should give additional doses of epinephrine if you need more than two injections for a single anaphylactic episode. Epinephrine Injection, USP Auto-Injector should **only** be injected into the middle of your outer thigh (upper leg), through clothing if necessary. Do not inject into your veins, buttocks, fingers, toes, hands or feet. Hold the leg of young children firmly in place before and during injection to prevent injuries. In case of accidental injection, please seek immediate medical treatment.

Rarely, patients who have used EpiPen[®] or EpiPen Jr[®] Auto-Injector may develop an infection at the injection site within a few days. Some of these infections can be serious. Call your healthcare professional right away if you have any of the following at an injection site: redness that does not go away, swelling, tenderness, or the area feels warm to the touch.

Tell your healthcare professional about all of your medical conditions, especially if you have asthma, a history of depression, thyroid problems, Parkinson's disease, diabetes, high blood pressure or heart problems, have any other medical conditions, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. Be sure to also tell your healthcare professional all the medicines you take, especially medicines for asthma. If you have certain medical conditions, or take certain medicines, your condition may get worse or you may have longer lasting side effects when you use EpiPen[®] or EpiPen Jr[®] Auto-Injector.

Common side effects include fast, irregular or "pounding" heartbeat, sweating, nausea or vomiting, breathing problems, paleness, dizziness, weakness, shakiness, headache, feelings of over excitement, nervousness or anxiety. These side effects usually go away quickly if you lie down and rest. **Tell your healthcare professional if you have any side effect that bothers you or that does not go away**.

Indications

EpiPen[®] and EpiPen Jr[®] Auto-Injectors are for the emergency treatment of life-threatening allergic reactions (anaphylaxis) caused by allergens, exercise, or unknown triggers; and for people who are at increased risk for these reactions. EpiPen[®] and EpiPen Jr[®] Auto-Injectors are intended for immediate administration as emergency supportive therapy only. Seek immediate emergency medical help right away.

Please see the full Prescribing Information and Patient Information

For additional information, please contact us at 800-395-3376.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit <u>www.fda.gov/medwatch</u> or call 1-800-FDA-1088.

EpiPen[®], EpiPen Jr[®], *EpiPen4Schools*[®] and the Mylan logo are registered trademarks of Mylan Inc.

© 2019 Mylan Specialty L.P. All rights reserved. EPI-2019-0021